

OXFORD CAMBRIDGE AND RSA EXAMINATIONS

**Advanced Subsidiary General Certificate of Education
Advanced General Certificate of Education**

MATHEMATICS

4724

Core Mathematics 4

Monday **23 JANUARY 2006** Afternoon 1 hour 30 minutes

Additional materials:
8 page answer booklet
Graph paper
List of Formulae (MF1)

TIME 1 hour 30 minutes

INSTRUCTIONS TO CANDIDATES

- Write your name, centre number and candidate number in the spaces provided on the answer booklet.
- Answer **all** the questions.
- Give non-exact numerical answers correct to 3 significant figures unless a different degree of accuracy is specified in the question or is clearly appropriate.
- You are permitted to use a graphical calculator in this paper.

INFORMATION FOR CANDIDATES

- The number of marks is given in brackets [] at the end of each question or part question.
- The total number of marks for this paper is 72.
- Questions carrying smaller numbers of marks are printed earlier in the paper, and questions carrying larger numbers of marks later in the paper.
- **You are reminded of the need for clear presentation in your answers.**

This question paper consists of 3 printed pages and 1 blank page.

- 1 Simplify $\frac{x^3 - 3x^2}{x^2 - 9}$. [3]
- 2 Given that $\sin y = xy + x^2$, find $\frac{dy}{dx}$ in terms of x and y . [5]
- 3 (i) Find the quotient and the remainder when $3x^3 - 2x^2 + x + 7$ is divided by $x^2 - 2x + 5$. [4]
 (ii) Hence, or otherwise, determine the values of the constants a and b such that, when $3x^3 - 2x^2 + ax + b$ is divided by $x^2 - 2x + 5$, there is no remainder. [2]
- 4 (i) Use integration by parts to find $\int x \sec^2 x \, dx$. [4]
 (ii) Hence find $\int x \tan^2 x \, dx$. [3]
- 5 A curve is given parametrically by the equations $x = t^2$, $y = 2t$.
 (i) Find $\frac{dy}{dx}$ in terms of t , giving your answer in its simplest form. [2]
 (ii) Show that the equation of the tangent to the curve at $(p^2, 2p)$ is

$$py = x + p^2. \quad [2]$$

 (iii) Find the coordinates of the point where the tangent at $(9, 6)$ meets the tangent at $(25, -10)$. [4]
- 6 (i) Show that the substitution $x = \sin^2 \theta$ transforms $\int \sqrt{\frac{x}{1-x}} \, dx$ to $\int 2 \sin^2 \theta \, d\theta$. [4]
 (ii) Hence find $\int_0^1 \sqrt{\frac{x}{1-x}} \, dx$. [5]
- 7 The expression $\frac{11 + 8x}{(2-x)(1+x)^2}$ is denoted by $f(x)$.
 (i) Express $f(x)$ in the form $\frac{A}{2-x} + \frac{B}{1+x} + \frac{C}{(1+x)^2}$, where A , B and C are constants. [5]
 (ii) Given that $|x| < 1$, find the first 3 terms in the expansion of $f(x)$ in ascending powers of x . [5]

- 8 (i) Solve the differential equation

$$\frac{dy}{dx} = \frac{2-x}{y-3},$$

giving the particular solution that satisfies the condition $y = 4$ when $x = 5$. [5]

- (ii) Show that this particular solution can be expressed in the form

$$(x-a)^2 + (y-b)^2 = k,$$

where the values of the constants a , b and k are to be stated. [3]

- (iii) Hence sketch the graph of the particular solution, indicating clearly its main features. [3]

- 9 Two lines have vector equations

$$\mathbf{r} = \begin{pmatrix} 4 \\ 2 \\ -6 \end{pmatrix} + t \begin{pmatrix} -8 \\ 1 \\ -2 \end{pmatrix} \quad \text{and} \quad \mathbf{r} = \begin{pmatrix} -2 \\ a \\ -2 \end{pmatrix} + s \begin{pmatrix} -9 \\ 2 \\ -5 \end{pmatrix},$$

where a is a constant.

- (i) Calculate the acute angle between the lines. [5]

- (ii) Given that these two lines intersect, find a and the point of intersection. [8]

BLANK PAGE

Permission to reproduce items where third-party owned material protected by copyright is included has been sought and cleared where possible. Every reasonable effort has been made by the publisher (OCR) to trace copyright holders, but if any items requiring clearance have unwittingly been included, the publisher will be pleased to make amends at the earliest possible opportunity.

OCR is part of the Cambridge Assessment Group. Cambridge Assessment is the brand name of University of Cambridge Local Examinations Syndicate (UCLES), which is itself a department of the University of Cambridge.